

Ulam-Hyers stability for partial differential inclusions

Vasile L. Lazăr

“Vasile Goldiș” Western University Arad, Romania.

M.Viteazul str., no.26, 440030, Satu Mare, Romania

E-mail: vasilazar@yahoo.com

Abstract

Using the weakly Picard operator technique, we will present Ulam-Hyers stability results for integral inclusions of Fredholm and Volterra type and for the Darboux problem associated to a partial differential inclusion.

2010 Mathematics Subject Classification: 47H10, 54H25, 54C60

Keywords and phrases: Ulam-Hyers stability, generalized Ulam-Hyers stability, multivalued weakly Picard operator, fixed point, selection, integral inclusion, partial differential inclusion.

1 Introduction

The Ulam stability (Ulam-Hyers, Ulam-Hyers-Rassias, Ulam-Hyers-Bourgin,...) of various functional equations has been investigated by many authors (see [14], [15], [6], [8], [3], [9], [13], [25], [30], [31]). There are

some results for differential equations ([16], [18], [19], [23], [36]), integral equations ([5], [17], [35]), for difference equations [4], [28], [29], [44]), etc. ([10], [11], [32]). For other results in the case of fixed point problems and coincidence point problems see [2], [26], [34], [37], [39].

The aim of this paper is to present existence and Ulam-Hyers stability results for some problems associated with integral inclusions and partial differential inclusions.

2 Ulam-Hyers stability via weakly Picard operators

Let (X, d) be a metric space and consider the following families of subsets of X :

$$P(X) := \{Y \in \mathcal{P}(X) \mid Y \neq \emptyset\}, \quad P_b(X) := \{Y \in P(X) \mid Y \text{ is bounded}\},$$

$$P_{cl}(X) := \{Y \in P(X) \mid Y \text{ is closed}\}, \quad P_{cp}(X) := \{Y \in P(X) \mid Y \text{ is compact}\}.$$

We will denote by $\bar{B}(x_0, r)$ the closure of $B(x_0, r)$ in (X, d) , where $B(x_0, r) := \{x \in X \mid d(x_0, x) < r\}$ is the open ball centered at $x_0 \in X$ with radius $r > 0$ and by $\tilde{B}(x_0, r)$ the closed ball centered at $x_0 \in X$ with radius $r > 0$, i.e., $\tilde{B}(x_0, r) := \{x \in X \mid d(x_0, x) \leq r\}$.

If (X, d) is a metric space, then the gap functional in $P(X)$ is defined as

$$D_d : P(X) \times P(X) \rightarrow \mathbb{R}_+, \quad D_d(A, B) = \inf\{d(a, b) \mid a \in A, b \in B\}.$$

In particular, if $x_0 \in X$ then $D_d(x_0, B) := D_d(\{x_0\}, B)$.

We will denote by H the generalized Pompeiu-Hausdorff functional on $P(X)$, defined as

$$H_d : P(X) \times P(X) \rightarrow \mathbb{R}_+ \cup \{+\infty\}, \quad H_d(A, B) = \max\{\sup_{a \in A} D_d(a, B), \sup_{b \in B} D_d(b, A)\}.$$

Let (X, d) be a metric space. If $F : X \rightarrow P(X)$ is a multivalued operator, then $x \in X$ is called a fixed point for F if and only if $x \in F(x)$. The

set $Fix(F) := \{x \in X \mid x \in F(x)\}$ is called the fixed point set of F , while $SFix(F) = \{x \in X \mid \{x\} = F(x)\}$ is called the strict fixed point set of F .

For a multivalued operator $F : X \rightarrow P(Y)$ the graph of F will be denoted by

$$Graph(F) := \{(x, y) \in X \times Y : y \in F(x)\}.$$

Notice that $f : X \rightarrow Y$ is a selection for $F : X \rightarrow P(Y)$ if $f(x) \in F(x)$, for each $x \in X$.

In particular, when F is a singlevalued operator, we obtain the similar well-known concepts in fixed point theory.

For the following notions see I.A. Rus [33] and [37], I.A. Rus, A. Petruşel, A. Sîntămărian [40] and A. Petruşel [27].

Definition 2.1. Let (X, d) be a metric space and $f : X \rightarrow X$ be an operator. By definition, f is a weakly Picard operator (briefly WPO) if the sequence $(f^n(x))_{n \in \mathbb{N}}$ of successive approximations for f starting from $x \in X$ converges, for all $x \in X$ and its limit is a fixed point of f .

If f is a WPO, then we consider the operator

$$f^\infty : X \rightarrow X \text{ defined by } f^\infty(x) := \lim_{n \rightarrow \infty} f^n(x).$$

Notice that $f^\infty(X) = Fix(f)$.

Definition 2.2. Let (X, d) be a metric space, $f : X \rightarrow X$ be a WPO and $c > 0$ be a real number. By definition, the operator f is a c -weakly Picard operator (briefly c -WPO) if and only if

$$d(x, f^\infty(x)) \leq c d(x, f(x)), \text{ for all } x \in X.$$

In the multivalued case we have the following concepts.

Definition 2.3. Let (X, d) be a metric space, and $F : X \rightarrow P_d(X)$ be a multivalued operator. By definition, F is a multivalued weakly Picard (briefly MWP) operator if for each $x \in X$ and each $y \in F(x)$ there exists a sequence

$(x_n)_{n \in \mathbb{N}}$ such that:

(i) $x_0 = x, x_1 = y$;

(ii) $x_{n+1} \in F(x_n)$, for each $n \in \mathbb{N}$;

(iii) the sequence $(x_n)_{n \in \mathbb{N}}$ is convergent and its limit is a fixed point of F .

Remark 2.1. A sequence $(x_n)_{n \in \mathbb{N}}$ satisfying condition (i) and (ii) in the Definition 2.3 is called a sequence of successive approximations of F starting from $(x, y) \in \text{Graph}(F)$.

If $F : X \rightarrow P(X)$ is a MWP operator, then we define $F^\infty : \text{Graph}(F) \rightarrow P(\text{Fix} F)$ by the formula $F^\infty(x, y) := \{ z \in \text{Fix}(F) \mid \text{there exists a sequence of successive approximations of } F \text{ starting from } (x, y) \text{ that converges to } z \}$.

Definition 2.4. Let (X, d) be a metric space and let $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be an increasing function which is continuous at 0 and $\psi(0) = 0$. Then $F : X \rightarrow P(X)$ is said to be a multivalued ψ -weakly Picard operator if it is a multivalued weakly Picard operator and there exists a selection $f^\infty : \text{Graph}(F) \rightarrow \text{Fix}(F)$ of F^∞ such that

$$d(x, f^\infty(x, y)) \leq \psi(d(x, y)), \quad \text{for all } (x, y) \in \text{Graph}(F).$$

If there exists $c > 0$ such that $\psi(t) = ct$, for each $t \in \mathbb{R}_+$, then F is called a multivalued c -weakly Picard operator.

Recall that, if (X, d) is a metric space, then $F : X \rightarrow P_{cl}(X)$ is said to be a multivalued α -contraction if $\alpha \in [0, 1)$ and

$$H_d(F(x), F(y)) \leq \alpha d(x, y), \quad \text{for all } x, y \in X,$$

Example 2.1. Let (X, d) be a complete metric space and $F : X \rightarrow P_{cl}(X)$ be a multivalued α -contraction. Then F is a c -MWP operator, where $c = (1 - \alpha)^{-1}$.

For the theory of weakly Picard operators, see [33] for the singlevalued case and [40] and [27] for the multivalued one.

We present now some Ulam-Hyers stability concepts for the fixed point problem associated with a multivalued operator.

Definition 2.5. Let (X, d) be a metric space and $F : X \rightarrow P(X)$ be a multivalued operator. The fixed point inclusion

$$(2.1) \quad x \in F(x), \quad x \in X$$

is called generalized Ulam-Hyers stable if and only if there exists $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ increasing, continuous at 0 and $\psi(0) = 0$ such that for each $\varepsilon > 0$ and for each solution $y^* \in X$ of the inequation

$$(2.2) \quad D_d(y, F(y)) \leq \varepsilon$$

there exists a solution x^* of the fixed point inclusion (2.1) such that

$$d(y^*, x^*) \leq \psi(\varepsilon).$$

If there exists $c > 0$ such that $\psi(t) := ct$, for each $t \in \mathbb{R}_+$, then the fixed point inclusion (2.1) is said to be Ulam-Hyers stable.

The following theorem is an abstract result concerning the Ulam-Hyers stability of the fixed point inclusion (2.1) for multivalued operators with compact values.

Theorem 2.1. (*I.A. Rus [37]*) *Let (X, d) be a metric space and $F : X \rightarrow P_{cp}(X)$ be a multivalued ψ -weakly Picard operator. Then, the fixed point inclusion (2.1) is generalized Ulam-Hyers stable.*

3 Existence and Ulam-Hyers stability for integral inclusions

We consider here some integral inclusion of Fredholm and Volterra type. Throughout this section we will denote by $\|\cdot\|$ the supremum norm in $C([a, b], \mathbb{R}^n)$ and by $|\cdot|$ a norm in \mathbb{R}^n .

Recall that $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is said to be a comparison function (see [38]) if it is increasing and $\varphi^k(t) \rightarrow 0$, as $k \rightarrow +\infty$. As a consequence, we also have $\varphi(t) < t$, for each $t > 0$, $\varphi(0) = 0$ and φ is continuous at 0.

Recall also the notion of strict comparison function. A function $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is said to be a strict comparison function (see [38]) if it is strictly increasing and $\sum_{n=1}^{\infty} \varphi^n(t) < +\infty$, for each $t > 0$.

The mappings $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ given by $\varphi(t) = at$ (where $a \in [0, 1[$) and respectively $\varphi(t) = \frac{t}{1+t}$, for each $t \in \mathbb{R}_+$ are examples of strict comparison functions.

The following result, a generalization of Covitz-Nadler fixed point principle (see [24], [7]) is known in the literature as Węgrzyk's fixed point theorem.

Theorem 3.2. *Let (X, d) be a complete metric space and $F : X \rightarrow P_{cl}(X)$ be a multivalued φ -contraction, i.e., $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a strict comparison function and*

$$H(F(x_1), F(x_2)) \leq \varphi(d(x_1, x_2)), \text{ for all } x_1, x_2 \in X.$$

Then $\text{Fix}(F)$ is nonempty and for any $x_0 \in X$ there exists a sequence of successive approximations of F starting from x_0 which converges to a fixed point of F .

Remark 3.2. *It is worth noting that, in the conditions of above result, if additionally $S\text{Fix}(F) \neq \emptyset$, then $\text{Fix}(F) = S\text{Fix}(F) = \{x^*\}$, see Sîntămărian [42]. Moreover, in this case, if the function $\beta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$, $\beta(t) := t - \varphi(t)$ is strictly increasing and onto, then, since*

$$d(x, x^*) \leq D(x, F(x)) + H(F(x), F(x^*)) \leq D(x, F(x)) + \varphi(d(x, x^*)), \text{ for all } x \in X,$$

we get that

$$d(x, x^*) \leq \beta^{-1}(D(x, F(x))), \text{ for all } x \in X,$$

This immediately implies that the fixed point problem $x \in F(x)$, $x \in X$ is generalized Ulam-Hyers stable with function β^{-1} .

Another Ulam-Hyers stability result, more efficient for applications, was proved in [21].

Theorem 3.3. Let (X, d) be a complete metric space and $F : X \rightarrow P_{cl}(X)$ be a multivalued φ -contraction. Then:

- (i) (existence of the fixed point) F is a MWP operator;
- (ii) (Ulam-Hyers stability for the fixed point inclusion) If additionally $\varphi(qt) \leq q\varphi(t)$ for every $t \in \mathbb{R}_+$ (where $q > 1$) and $t = 0$ is a point of uniform convergence for the series $\sum_{n=1}^{\infty} \varphi^n(t)$, then F is a ψ -MWP operator, with $\psi(t) := t + s(t)$, for each $t \in \mathbb{R}_+$ (where $s(t) := \sum_{n=1}^{\infty} \varphi^n(t)$);
- (iii) (data dependence of the fixed point set) Let $S : X \rightarrow P_{cl}(X)$ be a multivalued φ -contraction and $\eta > 0$ be such that $H(S(x), F(x)) \leq \eta$, for each $x \in X$. Suppose that $\varphi(qt) \leq q\varphi(t)$ for every $t \in \mathbb{R}_+$ (where $q > 1$) and $t = 0$ is a point of uniform convergence for the series $\sum_{n=1}^{\infty} \varphi^n(t)$. Then $H(Fix(S), Fix(F)) \leq \psi(\eta)$.

We will present now, using the above mentioned results, some existence and Ulam-Hyers stability theorems for multivalued operatorial inclusions.

Consider first the following Fredholm type integral inclusion.

$$(3.3) \quad x(t) \in \int_a^b K(t, s, x(s))ds + g(t), \quad t \in [a, b].$$

The main result concerning the stability of the Fredholm integral inclusion (3.3) is the following.

Theorem 3.4. Let $K : [a, b] \times [a, b] \times \mathbb{R}^n \rightarrow P_{cl,cv}(\mathbb{R}^n)$ and $g : [a, b] \rightarrow \mathbb{R}^n$ such that:

- (a) there exists an integrable function $M : [a, b] \rightarrow \mathbb{R}_+$ such that for each $t \in [a, b]$ and $u \in \mathbb{R}^n$ we have $K(t, s, u) \subset M(s)B(0; 1)$, a.e. $s \in [a, b]$;
- (b) for each $u \in \mathbb{R}^n$ $K(\cdot, \cdot, u) : [a, b] \times [a, b] \rightarrow P_{cl,cv}(\mathbb{R}^n)$ is jointly measurable;
- (c) for each $(s, u) \in [a, b] \times \mathbb{R}^n$ $K(\cdot, s, u) : [a, b] \rightarrow P_{cl,cv}(\mathbb{R}^n)$ is lower semi-continuous;

(d) there exists a continuous function $p : [a, b] \times [a, b] \rightarrow \mathbb{R}_+$ with $\sup_{t \in [a, b]} \int_a^b p(t, s) ds \leq 1$ and a strict comparison function $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that for each $(t, s) \in [a, b] \times [a, b]$ and each $u, v \in \mathbb{R}^n$ we have that

$$(3.4) \quad H(K(t, s, u), K(t, s, v)) \leq p(t, s) \cdot \varphi(|u - v|);$$

(e) g is continuous.

Then the following conclusions hold:

(a) the integral inclusion (3.3) has least one solution, i.e., there exists $x^* \in C([a, b], \mathbb{R}^n)$ which satisfies (3.3), for each $t \in [a, b]$.

(b) If additionally $\varphi(qt) \leq q\varphi(t)$ for every $t \in \mathbb{R}_+$ (where $q > 1$) and $t = 0$ is a point of uniform convergence for the series $\sum_{n=1}^{\infty} \varphi^n(t)$, then the integral inclusion (3.3) is generalized Ulam-Hyers stable with function ψ (where $\psi(t) := t + s(t)$, for each $t \in \mathbb{R}_+$ and $s(t) := \sum_{n=1}^{\infty} \varphi^n(t)$), i.e., for each $\varepsilon > 0$ and for any ε -solution y of (3.3), that is any $y \in C([a, b], \mathbb{R}^n)$ for which there exists $u \in C([a, b], \mathbb{R}^n)$ such that

$$u(t) \in \int_a^b K(t, s, y(s)) ds + g(t), \quad t \in [a, b]$$

and

$$|u(t) - y(t)| \leq \varepsilon, \quad \text{for each } t \in [a, b],$$

there exists a solution x^* of the integral inclusion (3.3) such that

$$|y(t) - x^*(t)| \leq \psi(\varepsilon), \quad \text{for each } t \in [a, b].$$

Moreover, in this case the continuous data dependence of the solution set of the integral inclusion (3.4) holds.

Proof. (a) Define the multivalued operator $T : C([a, b], \mathbb{R}^n) \rightarrow \mathcal{P}(C([a, b], \mathbb{R}^n))$ by

$$T(x) := \left\{ v \in C([a, b], \mathbb{R}^n) \mid v(t) \in \int_a^b K(t, s, x(s)) ds + g(t), \quad t \in [a, b] \right\}.$$

Then, (3.3) is equivalent to the fixed point inclusion

$$(3.5) \quad x \in T(x), \quad x \in C([a, b], \mathbb{R}^n).$$

The proof is organized in several steps. We successively prove:

$$1. T(x) \in P_{cp}(C([a, b], \mathbb{R}^n)).$$

From (e) and Theorem 2 in Rybiński [41] we have that for each $x \in C([a, b], \mathbb{R}^n)$ there exists $k(t, s) \in K(t, s, x(s))$, for all $(t, s) \in [a, b]$, such that $k(t, s)$ is integrable with respect to s and continuous with respect to t . Then $v(t) := \int_a^b k(t, s)ds + g(t)$, has the property $v \in T(x)$. Moreover, from (a) and (b), via Theorem 8.6.3. in Aubin and Frankowska [1], we get that $T(x)$ is a compact set, for each $x \in C([a, b], \mathbb{R}^n)$.

$$2. H(T(x_1), T(x_2)) \leq \varphi(\|x_1 - x_2\|), \text{ for each } x_1, x_2 \in C([a, b], \mathbb{R}^n).$$

Notice first that one may suppose (without affecting the generality of the Lipschitz condition) that the inequality (3.4) is strict. Let $x_1, x_2 \in C([a, b], \mathbb{R}^n)$ and $v_1 \in T(x_1)$. Then $v_1(t) \in \int_a^b K(t, s, x_1(s))ds + g(t)$, $t \in [a, b]$. It follows that $v_1(t) = \int_a^b k_1(t, s)ds + g(t)$, $t \in [a, b]$, for some $k_1(t, s) \in K(t, s, x_1(s))$, $(t, s) \in [a, b] \times [a, b]$.

From (d) we have $H(K(t, s, x_1(s)), K(t, s, x_2(s))) < p(t, s)\varphi(|x_1(s) - x_2(s)|) \leq p(t, s)\varphi(\|x_1 - x_2\|)$. Thus, there exists $w \in K(t, s, x_2(s))$ such that $|k_1(t, s) - w| \leq p(t, s)\varphi(\|x_1 - x_2\|)$, for $t, s \in [a, b]$.

Let us define $U : [a, b] \times [a, b] \rightarrow P(\mathbb{R}^n)$, by $U(t, s) = \{w \mid |k_1(t, s) - w| \leq p(t, s)\varphi(\|x_1 - x_2\|)\}$. Since the multi-valued operator $V(t, s) := U(t, s) \cap K(t, s, x_2(s))$ is jointly measurable and lower semi-continuous in t there exists $k_2(t, s)$ a selection for V , jointly measurable (and, hence, integrable in s) and continuous in t . Hence, $k_2(t, s) \in K(t, s, x_2(s))$ and $|k_1(t, s) - k_2(t, s)| \leq p(t, s)\varphi(\|x_1 - x_2\|)$, for each $t, s \in [a, b]$.

Consider $v_2(t) = \int_a^b k_2(t, s)ds + g(t)$, $t \in [a, b]$. Then, we have:

$$|v_1(t) - v_2(t)| \leq \int_a^b |k_1(t, s) - k_2(t, s)| ds \leq \int_a^b p(t, s) \varphi(\|x_1 - x_2\|) ds \leq \varphi(\|x_1 - x_2\|).$$

A similar relation can be obtained by interchanging the roles of x_1 and x_2 . Thus the second step follows.

The first conclusion follows by the above mentioned Węgrzyk's fixed point theorem, see Theorem 3.3 (i) (see also [43]).

(b) We will prove that the fixed point inclusion problem (3.5) is generalized Ulam-Hyers stable. Indeed, let $\varepsilon > 0$ and $y \in C([a, b], \mathbb{R}^n)$ for which there exists $u \in C([a, b], \mathbb{R}^n)$ such that

$$u(t) \in \int_a^b K(t, s, y(s)) ds + g(t), \quad t \in [a, b]$$

$$\text{and } \|u - y\| \leq \varepsilon.$$

Then $D_{\|\cdot\|}(y, T(y)) \leq \varepsilon$. Moreover, by the above proof we have that T is a multivalued φ -contraction and using Theorem 3.3(i)-(ii), we obtain that T is a multivalued ψ -weakly Picard operator. Then, by Theorem 2.1 we obtain that the fixed point problem (3.5) is generalized Ulam-Hyers stable. Thus, the integral inclusion (3.4) is generalized Ulam-Hyers stable.

Concerning the last conclusion of the theorem, we apply Theorem 3.3 (iii). □

A second application concerns an integral inclusion of Volterra type.

$$(3.6) \quad x(t) \in \int_a^t K(t, s, x(s)) ds + g(t), \quad t \in [a, b].$$

By a similar method, we can prove the following.

Theorem 3.5. *Let $K : [a, b] \times [a, b] \times \mathbb{R}^n \rightarrow P_{cl,cv}(\mathbb{R}^n)$ and $g : [a, b] \rightarrow \mathbb{R}^n$ such that:*

(a) there exists an integrable function $M : [a, b] \rightarrow \mathbb{R}_+$ such that for each $t \in [a, b]$ and $u \in \mathbb{R}^n$ we have $K(t, s, u) \subset M(s)B(0; 1)$, a.e. $s \in [a, b]$;

(b) for each $u \in \mathbb{R}^n$ $K(\cdot, \cdot, u) : [a, b] \times [a, b] \rightarrow P_{cl,cv}(\mathbb{R}^n)$ is jointly measurable;

(c) for each $(s, u) \in [a, b] \times \mathbb{R}^n$ $K(\cdot, s, u) : [a, b] \rightarrow P_{cl,cv}(\mathbb{R}^n)$ is lower semi-continuous;

(d) there exists a continuous function $p : [a, b] \rightarrow \mathbb{R}_+^*$ and a strict comparison function $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\varphi(\lambda t) \leq \lambda \varphi(t)$, for each $t \in \mathbb{R}_+$ and each $\lambda \geq 1$, such that for each $(t, s) \in [a, b] \times [a, b]$ and each $u, v \in \mathbb{R}^n$ we have that

$$(3.7) \quad H(K(t, s, u), K(t, s, v)) \leq p(s) \cdot \varphi(|u - v|);$$

(e) g is continuous.

Then the following conclusions hold:

(a) the integral inclusion (3.6) has at least one solution, i.e., there exists $x^* \in C([a, b], \mathbb{R}^n)$ which satisfies (3.6) for each $t \in [a, b]$;

(b) If additionally $\varphi(qt) \leq q\varphi(t)$ for every $t \in \mathbb{R}_+$ (where $q > 1$) and $t = 0$ is a point of uniform convergence for the series $\sum_{n=1}^{\infty} \varphi^n(t)$, then the integral inclusion (3.3) is generalized Ulam-Hyers stable with function ψ (where $\psi(t) := t + s(t)$, for each $t \in \mathbb{R}_+$ and $s(t) := \sum_{n=1}^{\infty} \varphi^n(t)$), i.e., for each $\varepsilon > 0$ and for any ε -solution y of (3.6), that is, any $y \in C([a, b], \mathbb{R}^n)$ for which there exists $u \in C([a, b], \mathbb{R}^n)$ such that

$$u(t) \in \int_a^t K(t, s, y(s))ds + g(t), \quad t \in [a, b]$$

and

$$|u(t) - y(t)| \leq \varepsilon, \quad \text{for each } t \in [a, b],$$

there exists a solution x^* of the integral inclusion (3.6) such that

$$|y(t) - x^*(t)| \leq \psi(c\varepsilon), \quad \text{for each } t \in [a, b] \text{ and some } c > 0.$$

Moreover, in this case the continuous data dependence of the solution set of the integral inclusion (3.7) holds.

Proof. We consider the multi-valued operator $T : C([a, b], \mathbb{R}^n) \rightarrow \mathcal{P}(C([a, b], \mathbb{R}^n))$

$$T(x) := \left\{ v \in C([a, b], \mathbb{R}^n) \mid v(t) \in \int_a^t K(t, s, x(s))ds + g(t), t \in [a, b] \right\}.$$

Then, (3.6) is equivalent to the fixed point inclusion

$$(3.8) \quad x \in T(x), \quad x \in C([a, b], \mathbb{R}^n).$$

As in the proof of Theorem 3.4 we obtain $T(x) \in P_{cp}(C([a, b], \mathbb{R}^n))$. Next, we will prove that T is a multivalued φ -contraction on $C([a, b], \mathbb{R}^n)$.

Notice first that one may suppose (without affecting the generality of the Lipschitz condition) that the inequality (3.7) is strict. Let $x_1, x_2 \in C([a, b], \mathbb{R}^n)$ and $v_1 \in T(x_1)$. Then $v_1(t) \in \int_a^t K(t, s, x_1(s))ds + g(t)$, $t \in [a, b]$. It follows that $v_1(t) = \int_a^b k_1(t, s)ds + g(t)$, $t \in [a, b]$, for some $k_1(t, s) \in K(t, s, x_1(s))$, $(t, s) \in [a, b] \times [a, b]$.

From (d) we have $H(K(t, s, x_1(s)), K(t, s, x_2(s))) < p(s)\varphi(|x_1(s) - x_2(s)|)$. Thus, there exists $w \in K(t, s, x_2(s))$ such that $|k_1(t, s) - w| \leq p(s)\varphi(|x_1(s) - x_2(s)|)$, for $t, s \in [a, b]$.

Let us define $U : [a, b] \times [a, b] \rightarrow P(\mathbb{R}^n)$, by $U(t, s) = \{w \mid |k_1(t, s) - w| \leq p(t, s)\varphi(|x_1(s) - x_2(s)|)\}$. Since the multivalued operator $V(t, s) := U(t, s) \cap K(t, s, x_2(s))$ is jointly measurable and lower semi-continuous in t there exists $k_2(t, s)$ a selection for V , jointly measurable (hence, integrable in s) and continuous in t . Hence, $k_2(t, s) \in K(t, s, x_2(s))$ and $|k_1(t, s) - k_2(t, s)| \leq p(s)\varphi(|x_1(s) - x_2(s)|)$, for each $t, s \in [a, b]$.

Consider $v_2(t) = \int_a^t k_2(t, s)ds + g(t)$, $t \in [a, b]$. We denote by $\|\cdot\|_B$ a Bielecki-type norm in $C([a, b], \mathbb{R}^n)$, given by $\|x\|_B := \sup_{t \in [a, b]} (|x(t)|e^{-q(t)})$, where $q(t) := \int_a^t p(s)ds$.

Then, for each $t \in [a, b]$, we have:

$|v_1(t) - v_2(t)| \leq \int_a^t |k_1(t, s) - k_2(t, s)| ds \leq \int_a^t p(s) \varphi(|x_1(s) - x_2(s)|) ds =$
 $\int_a^t p(s) \varphi(e^{q(s)} |x_1(s) - x_2(s)| e^{q(s)}) ds \leq \int_a^t p(s) e^{q(s)} \varphi(\|x_1 - x_2\|_B) ds =$
 $\varphi(\|x_1 - x_2\|_B) (e^{q(t)} - e^{q(a)}) \leq \varphi(\|x_1 - x_2\|_B) e^{q(t)}.$ Thus, we immediately get

$$\|v_1 - v_2\|_B \leq \varphi(\|x_1 - x_2\|_B).$$

A similar relation can be obtained by interchanging the roles of x_1 and x_2 . Thus, we have that

$$H_{\|\cdot\|_B}(T(x_1), T(x_2)) \leq \varphi(\|x_1 - x_2\|_B), \text{ for each } x_1, x_2 \in C([a, b], \mathbb{R}^n),$$

which proves that T is a multivalued φ -contraction. The conclusion (a) follows by the above mentioned Węgrzyk's fixed point theorem, see Theorem 3.3 (i) (see also [43]).

(b) We will prove that the fixed point inclusion problem (3.6) is generalized Ulam-Hyers stable. For this purpose, it is enough to prove that the fixed point inclusion problem (3.8) is generalized Ulam-Hyers stable. For this purpose, let $\varepsilon > 0$ and $y \in C([a, b], \mathbb{R}^n)$ for which there exists $u \in C([a, b], \mathbb{R}^n)$ such that

$$u(t) \in \int_a^t K(t, s, y(s)) ds + g(t), \quad t \in [a, b]$$

and

$$|u(t) - y(t)| \leq \varepsilon, \text{ for each } t \in [a, b].$$

Notice that

$$\|\cdot\|_B \leq \|\cdot\| \leq \|\cdot\|_B e^{\tau q(b)}.$$

Then, we obtain that $\|u - y\|_B \leq \|u - y\| \leq \varepsilon$. Thus, $D_{\|\cdot\|_B}(y, T(y)) \leq \varepsilon$. Moreover, by the above proof, T is a multivalued φ -contraction with respect to $\|\cdot\|_B$ and, thus, T is a MWP operator. Using Theorem 3.3(i)-(ii), we obtain that T is a multivalued ψ -MWP operator. Thus, conclusion (b) is a consequence of Theorem 2.1. Hence, there exists a solution x^* of the integral inclusion (3.6) such that

$$\|y - x^*\|_B \leq \psi(\varepsilon).$$

Hence,

$$|y(t) - x^*(t)| \leq \psi(e^{\tau q(b)} \varepsilon), \text{ for each } t \in [a, b].$$

Concerning the last conclusion of the theorem, we apply Theorem 3.3 (iii).

□

4 Existence and Ulam-Hyers stability for partial differential inclusions

Let us consider the following Darboux problem for a second order differential inclusion

$$(4.9) \quad \begin{cases} \frac{\partial^2 u}{\partial x \partial y} \in F(x, y, u(x, y)) \\ u(x, 0) = \lambda(x, 0), \quad u(0, y) = \lambda(0, y), \end{cases}$$

where $F : I_1 \times I_2 \times \mathbb{R}^m \rightarrow P_{cl}(\mathbb{R}^m)$ (with $I_i = [0, T_i]$, $i \in \{1, 2\}$) and $\lambda(x, y) = \alpha(x) + \beta(y) - \alpha(0)$ (with α, β continuous functions on I_1 respectively I_2 and $\alpha(0) = \beta(0)$).

Denote by $\Pi = I_1 \times I_2$ and let $a > 0$. By L^1 we will denote the Banach space of all measurable Lebesgue functions $\eta : \Pi \rightarrow \mathbb{R}^m$, endowed with the norm

$$\|\eta\|_1 = \int_{\Pi} \int e^{-a(x+y)} |\eta(x, y)| dx dy.$$

Let C be the Banach space of continuous functions $u : \Pi \rightarrow \mathbb{R}^m$, with the norm $\|u\|_C = \sup_{(x,y) \in \Pi} |u(x, y)|$ and let \tilde{C} be the linear subspace of C consisting of all $\lambda \in C$ such that there exist continuous functions $\alpha \in C(I_1, \mathbb{R}^m)$ and $\beta \in C(I_2, \mathbb{R}^m)$ with $\alpha(0) = \beta(0)$ satisfying $\lambda(x, y) = \alpha(x) + \beta(y) - \alpha(0)$, for all $x, y \in I_1 \times I_2$. Obviously, \tilde{C} with the norm of C is a separable Banach space.

By definition, the Darboux problem (4.9) is called Ulam-Hyers stable if for each $\varepsilon > 0$ and for any ε -solution w of (4.9), there exists a solution u^* of (4.9) such that $|w(x, y) - u^*(x, y)| \leq c\varepsilon$, for each $(x, y) \in \Pi$ and for some $c > 0$.

We have the following existence and Ulam-Hyers stability result.

Theorem 4.6. *Consider the Darboux Problem (4.9) and suppose that the above mentioned conditions hold. Suppose also that the following assumptions hold:*

- i) for each $u \in \mathbb{R}^m$, $F(\cdot, \cdot, u)$ is measurable;*
- ii) there exists $k > 0$ such that a.e. $(x, y) \in I_1 \times I_2$ the multifunction $F(x, y, \cdot)$ is k -Lipschitz;*
- iii) $a > \sqrt{k}$.*

Then, the Darboux Problem (4.9) has at least one solution and it is Ulam-Hyers stable.

Proof. For $\lambda \in \tilde{C}, \eta \in L^1$ define

$$T_\lambda(\eta) := \{\mu \in L^1 : \mu(x, y) \in M_{\lambda, \eta}(x, y), \text{ a. e. on } \Pi\},$$

where

$$M_{\lambda, \eta}(x, y) = F(x, y, \lambda(x, y) + \int_0^x \int_0^y \eta(s, t) ds dt), (s, t) \in \Pi.$$

Notice that F_{T_λ} coincides with the solution set of the considered problem. Moreover, we have that $T_\lambda : L^1 \rightarrow P_{cl}(L^1)$ and it is a MWP operator. Indeed, we have

$$H_1(T_\lambda(\eta_1), T_\lambda(\eta_2)) \leq \frac{k}{a^2} \cdot \|\eta_1 - \eta_2\|_1, \text{ for all } \lambda \in \tilde{C} \text{ and } \eta_1, \eta_2 \in L^1.$$

Thus, T_λ is a $\frac{k}{a^2}$ -multivalued contraction on L^1 and hence is a MWP operator. Thus, there exists $u^* \in L^1$ a fixed point for T_λ , which is also a solution for the Darboux Problem (4.9). For the second part of our theorem it is enough to prove that T_λ is a multivalued c -weakly Picard operator. Since T_λ is a $\frac{k}{a^2}$ -multivalued contraction on L^1 , we immediately get (see Example 2.1) that T_λ is a multivalued c -weakly Picard operator with $c := \frac{1}{1 - ka^{-2}}$. Thus, the second conclusion follows by Theorem 2.1. \square

References

- [1] J.-P. Aubin, H. Frankowska, *Set-Valued Analysis*, Birkhauser, Basel, 1990.
- [2] M.F. Bota-Boriceanu, A. Petruşel, *Ulam-Hyers stability for operatorial equations*, Analele Univ. Al. I. Cuza Iaşi, Math., 2011, to appear.
- [3] W.W. Breckner, and T. Trif, *Convex Functions and Related Functional Equations*, Cluj Univ. Press, Cluj-Napoca, 2008.
- [4] J. Brzdek, D. Popa, B. Xu, *The Hyers-Ulam stability of nonlinear recurrences*, J. Math. Anal. Appl., **335**(2007), 443-449.
- [5] L.P. Castro, A. Ramos, Hyers-Ulam-Rassias stability for a class of Volterra integral equations, *Banach J. Math. Anal.*, 3(2009), no.1, 36-43.
- [6] L. Cădariu, *Stabilitate Ulam-Hyers-Bourgin pentru ecuaţii funcţionale*, Ed. Univ. de Vest, Timişoara, 2007.
- [7] H. Covitz, S.B. Nadler jr., Multivalued contraction mappings in generalized metric spaces, *Israel J. Math.*, 8(1970), 5-11.
- [8] S. Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, 2002.
- [9] P. Găvruta, *On a problem of G. Isac and Th. Rassias concerning the stability of mapping*, J. Math. Anal. Appl., **261**(2001), 543-553.
- [10] P.M. Gruber, *Stability of isometries*, Trans. Amer. Math. Soc., **245**(1978), 263-277.
- [11] G. Hirisawa, T. Miura, *Hyers-Ulam stability of a closed operator in a Hilbert space*, Bull. Korean Math. Soc., **43**(2006), No. 1, 107-117.
- [12] S. Hu, N.S. Papageorgiou, *Handbook of Multivalued Analysis, Vol. I and II*, Kluwer Acad. Publ., Dordrecht, 1997, 1999.

- [13] D.H. Hyers, *The stability of homomorphism and related topics*, in: Global Analysis-Analysis on Manifolds (Th.M. Rassias, ed.), Teubner, Leipzig, 1983, 140-153.
- [14] D.H. Hyers, G. Isac, Th.M. Rassias, *Stability of Functional Equations in Several Variables*, Birkhäuser, Basel, 1998.
- [15] S.-M. Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press, Palm Harbor, 2001.
- [16] S.-M. Jung, *Hyers-Ulam stability of first order linear differential equations with constant coefficients*, J. Math. Anal. Appl., **320**(2006), 549-561.
- [17] S.-M. Jung, *A fixed point approach to the stability of a Volterra integral equation*, Fixed Point Theory and Applications, Vol. 2007 (2007), Article ID 57064, 1-9, doi:10.1155/2007/57064.
- [18] S.-M. Jung, K.-S. Lee, *Hyers-Ulam stability of first order linear partial differential equations with constant coefficients*, Math. Ineq. Appl., **10**(2007), no. 2, 261-266.
- [19] S.-M. Jung, Th.M. Rassias, *Generalized Hyers-Ulam stability of Riccati differential equation*, Math. Ineq. Appl., **11**(2008), No. 4, 777-782.
- [20] T.A. Lazăr, A. Petruşel, N. Shahzad, Fixed points for non-self operators and domain invariance theorems, *Nonlinear Anal.*, 70 (2009), 117-125.
- [21] V.L. Lazăr, Fixed point theory for multivalued φ -contractions, *Fixed Point Theory and Applications*, vol. 2011:50, (2011), 1-12, doi:10.1186/1687-1812-2011-50.
- [22] J. Matkowski, Integrable solutions of functional equations, *Dissertationes Math. (Rozprawy Mat.)* **127**(1975) 68 pp.

- [23] T. Miura, S.-M. Jung, S.-E. Takahasi, *Hyers-Ulam-Rassias stability of the Banach space valued linear differential equation $y' = \lambda y$* , J. Korean Math. Soc., **41**(2004), 995-1005.
- [24] S.B. Nadler jr., Multivalued contraction mappings, *Pacific J. Math.*, 30(1969), 475-488.
- [25] Zs. Páles, *Generalized stability of the Cauchy functional equation*, Aequationes Math., **56**(1998), no. 3, 222-232.
- [26] T.P. Petru, A. Petruşel, J.-C. Yao, *Ulam-Hyers stability for operatorial equations and inclusions via nonself operators*, Taiwanese J. Math., 2011, to appear.
- [27] A. Petruşel, *Multivalued weakly Picard operators and applications*, Sci. Math. Japon., **59**(2004), 167-202.
- [28] D. Popa, *Hyers-Ulam-Rassias stability of linear recurrence*, J. Math. Anal. Appl., **309**(2005), 591-597.
- [29] D. Popa, *Hyers-Ulam stability of the linear recurrence with constant coefficients*, Advances in Difference Equations, **2**(2005), 101-107.
- [30] V. Radu, *The fixed point alternative and the stability of functional equations*, Fixed Point Theory, **4**(2003), No. 1, 91-96.
- [31] Th. M. Rassias, *On the stability of the linear mappings in Banach spaces*, Proc. Amer. Math. Soc., **72**(1978), 297-300.
- [32] S. Reich, A.J. Zaslavski, *A stability result in fixed point theory*, Fixed Point Theory, **6**(2005), No. 1, 113-118.
- [33] I.A. Rus, *Picard operators and applications*, Scientiae Mathematicae Japonicae, **58**(2003), No. 1, 191-219.

- [34] I.A. Rus, *The theory of a metrical fixed point theorem: theoretical and applicative relevances*, Fixed Point Theory, **9**(2008), No. 2, 541-559.
- [35] I.A. Rus, *Gronwall lemma approach to the Hyers-Ulam-Rassias stability of an integral equation*, in: Nonlinear Analysis and Variational Problems, pp. 147-152 (P. Pardalos, Th. M. Rassias and A.A. Khan (Eds.)), Springer, 2009.
- [36] I.A. Rus, *Ulam stability of ordinary differential equations*, Studia Univ. Babeş-Bolyai, Math., **54**(2009), No. 4, 125-133.
- [37] I.A. Rus, *Remarks on Ulam stability of the operatorial equations*, Fixed Point Theory, **10**(2009), No. 2, 305-320.
- [38] I.A. Rus, Generalized Contractions and Applications, Cluj University Press, Cluj-Napoca, 2001.
- [39] I.A. Rus, A. Petruşel, G. Petruşel, *Fixed Point Theory*, Cluj University Press, 2008.
- [40] I.A. Rus, A. Petruşel, A. Sîntămărian, Data dependence of the fixed points set of some multivalued weakly Picard operators, *Nonlinear Anal.*, 2003, 52, 1947-1959.
- [41] L. Rybinski, On Carathéodory type selections, *Fund. Math.*, 125(1985), 187-193.
- [42] A. Sîntămărian, Data dependence of the fixed points of some Picard operators. (English) *Semin. Fixed Point Theory Cluj-Napoca* 2(2001), 81-86.
- [43] R. Węgrzyk, Fixed point theorems for multifunctions and their applications to functional equations, *Dissertationes Math.*, 201(1982), 28 pp.
- [44] M. Xu, *Hyers-Ulam-Rassias stability of a system of first order linear recurrences*, Bull. Korean Math. Soc., **44**(2007), No. 4, 841-849.

(Received December 1, 2011)